

# 南極海跡湖ぬるめ池における原生生物マイクロプランクトンの群集構造解析

久保伸一朗<sup>1</sup>、茶屋 葵<sup>1</sup>、川又明德<sup>2</sup>、小杉真貴子<sup>3</sup>、伊村智<sup>3</sup>、黒沢則夫<sup>1</sup>

<sup>1</sup>創価大学

<sup>2</sup>愛媛県総合科学博物館

<sup>3</sup>国立極地研究所

## Community analysis of protistan microplankton of Antarctic marine relic lake Nurume-Ike

Shinichiro Kubo<sup>1</sup>, Aoi Chaya<sup>1</sup>, Akinori Kawamata<sup>2</sup>, Makiko Kosugi<sup>3</sup>, Satoshi Imura<sup>3</sup>, and Norio Kurosawa<sup>1</sup>

<sup>1</sup>Soka University

<sup>2</sup>Ehime Prefectural Science Museum

<sup>3</sup>National Institute of Polar Research

Protists are essential to the function of most aquatic ecosystems and play pivotal roles as primary producers and consumers (Azam *et al.* 1983; Caron *et al.* 1999). It is important to clarify the diversity and community structure of protists to understand the aquatic ecosystem in an area of interest. In this study, we attempted to reveal the biodiversity and community structure of protistan microplankton in Antarctic lake Nurume-Ike. Nurume-Ike, located in Langhovde area, is a typical marine relic and meromictic lake (Sano *et al.*, 1977). The water column of this lake consists of three to four layers (Kudoh *et al.*, 2008) as follows. The surface layer (0-3 m) is slightly saline and is influenced by melt ice and water input from the surrounding catchment. The salinity of the second and third layers (3-11 m) is similar to that of the ocean. The bottom layer (>11 m) has one and a half times the salinity of the ocean, and has a completely reduced environment.

Sampling was conducted at Nurume-Ike (69° 13' 23" S, 39° 39' 43" E) on January 3, 2013 during the 54th Japanese Antarctic Research Expedition. The water temperature and pH were 9.3°C and 8.1, respectively. Ten liters of surface water was screened through a 100-μm nylon mesh and poured into a container. The cells were collected on membrane filters (pore size 10 μm), fixed with 5% Lugol's solution, and stored at 0-4°C until analysis. The fixed cells were recovered by centrifugation, and metagenomic DNA was extracted. The protistan 18S rRNA gene clone library was constructed using the metagenomic DNA. The clones were randomly chosen, sequenced, and identified by using BLAST (Altschul *et al.* 1990).

The protistan microplankton in Nurume-Ike was not too diverse compared to ordinally aquatic environments in temperate regions. Total 146 clones were grouped into 11 phylotypes. Most dominant phylotype, NRM-009, showed 99% sequence similarity with nonmotile dinoflagellate *Spiniferodinium galeiforme* (Horiguchi and Chihara, 1987) originally isolated in subtropical island in Japan. This phylotype shared 67% of total clones. The second dominant phylotype, NRM-082, was related to Prasinophycean flagellate *Pyramimonas gelidicola* (McFadden *et al.*, 1982) originally isolated from sea ice in Prydz Bay, East Antarctica. The phylogenetic and taxonomic analyses of other phylotypes are now investigated. It will be also interesting to see the genetic differences between Nurume-Ike protistan species and their marine relatives as topics of evolutionary biology.

## References

- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol.* 215:403-410.
- Azam F, Fenchel T, Field JG, Gray JS, Meyer-Reil LA, Thingstad F (1983) The ecological role of water-column microbes in the sea. *Mar Ecol Prog Ser.* 10:257-263.
- Caron DA, Lim EL, Dennett MR, Gast RJ, Kosman C, Delong EF (1999) Molecular phylogenetic analysis of the heterotrophic chrysophyte genus *Paraphysomonas* (Chrysophyceae), and the design of rRNA-targeted oligonucleotide probes for two species. *J Phycol.* 35:824-837.
- Horiguchi T, Chihara M (1987) *Spiniferodinium galeiforme*, a new genus and species of benthic dinoflagellates (Phytodiniales, Pyrrhophyta) from Japan. *Phycologia.* 26:478-487.
- Kudoh S, Tanabe Y, Takahashi K (2008) Abundance of benthic copepods in a saline lake in East Antarctica. *Polar Biol.* 31:1539-1542.
- McFadden GI, Moestrup Ø, Wetherbee R (1982) *Pyramimonas gelidicola* sp. nov. (Prasinophyceae), a new species isolated from Antarctic sea ice. *Phycologia.* 21:103-111.
- Sano M, Nakai N, Torii T (1977) On the two-layer structure in Lake Nurume, Antarctica. *Antarc Res* 58:63-68 (in Japanese with English abstract).